

F4



(11) Publication number: **0 424 059 A1**

(12) **EUROPEAN PATENT APPLICATION**

(21) Application number: 90311245.6

(51) Int. Cl.⁵: **A61K 31/415**

(22) Date of filing: 15.10.90

(30) Priority: 17.10.89 FI 894911

(43) Date of publication of application:
24.04.91 Bulletin 91/17

(84) Designated Contracting States:
AT BE CH DE DK FR GB IT LI LU NL SE


(71) Applicant: Orion-yhtymä Oy
Orionintie 1, PL 65
SF-02101 Espoo(FI)

(72) Inventor: Kalso, Elja
Aarnivalkeantie 5 A 15
SF-02100 Espoo(FI)
Inventor: Lammintausta, Risto
Meltolistentie
SF-20900 Turku(FI)

(74) Representative: Sexton, Jane Helen et al
J.A. Kemp & Co. 14 South Square Gray's Inn
London WC1R 5LX(GB)

(54) Use of substituted imidazoles.

(57) This invention relates to local epidural or intraspinal use of compounds of the formula:




where X is H or CH₃ and R₁ and R₂, which can be the same or different, are each H or CH₃ and their stereoisomers and their non-toxic, pharmaceutically acceptable salts. Especially useful are the compounds known under the generic names medetomidine ((±)-4-[1-(2,3-dimethylphenyl)ethyl]-1H-imidazole), dex-medetomidine (+)-4-[1-(2,3-dimethylphenyl)ethyl]-1H-imidazole and detomidine 4-(2,3-dimethylbenzyl)-1H-imidazole.

EP 0 424 059 A1

USE OF SUBSTITUTED IMIDAZOLES

This invention relates to local epidural or intraspinal use of compounds which are imidazoles of the formula



where X is H or CH₃ and R₁ and R₂, which can be the same or different, are each H or CH₃, their stereoisomers and their non-toxic, pharmaceutically acceptable salts. Especially useful are the compounds known under the generic names medetomidine ((±)-4-[1-(2,3-dimethylphenyl)ethyl]-1H-imidazole, dex-
 15 medetomidine ((+)-4-[1-(2,3-dimethylphenyl)ethyl]-1H-imidazole and detomidine (4-(2,3-dimethylbenzyl)-1H-imidazole.


Compounds of formula (I), particularly dexmedetomidine, medetomidine and detomidine are potent and selective α₂-adrenoceptor agonists. Using systemic administration they have been shown to be potent sedative, hypotensive, and analgesic compounds and useful in anxiolytic and perioperative treatment. These
 20 compounds and uses have been described in earlier publications e.g. European Patent Publications 24829, 72615, 187471, 270267, 300652 and 331374. The usefulness of the compounds as a painkiller for chronic use is, however, limited since analgesic effects are achieved in conjunction with α₂-adrenoceptor mediated pharmacological effects, including hypotension and sedation.

Intravenous administration of 100-300 μg/kg of the compound results in an analgesic effect, but also
 25 has a sedative and a hypotensive effect.

Intravenous administration of 1-30 μg/kg results in an anxiolytic effect but no appreciable analgesia.


The applicants have now identified a method of obtaining analgesia without producing a sedative or hypotensive effect. Such analgesia may be obtained by administering the compound intrathecally i.e. by a
 30 local epidural or intraspinal route.

Thus the present invention provides the use of a compound which is an imidazole of the formula (I):



40 where X is H or CH₃ and R₁ and R₂, which can be the same or different, are each H or CH₃, a stereoisomer thereof or a non-toxic pharmaceutically acceptable salt thereof in the manufacture of a medicament for local epidural or intraspinal administration.

The present invention also provides a method for obtaining analgesia in a mammal comprising local epidural or intraspinal administration to the mammal of a compound which is an imidazole of the formula (I):
 45



where X is H or CH₃ and R₁ and R₂, which can be the same or different, are H or CH₃, a stereoisomer

thereof or a nontoxic, pharmaceutically acceptable salt thereof, in an amount effective to achieve the desired level of analgesia.

Using this method it is only necessary to administer the compound in an amount which, if administered systemically e.g. intravenously, would typically produce an anxiolytic effect. For example, dexmedetomidine administered to rats produced an almost complete antinociceptive effect at very low intrathecal doses of compound e.g. 1-30 $\mu\text{g/kg}$ for rats using a standard analgesia model, the tail-flick test (ref: D'Amour F.E., Smith D.L: A Method For Determining Loss of Pain Sensation. J. Pharmacol. Exp. Ther. 1941, 72: 74-79).

To achieve a similar antinociceptive effect using intravenous administration typically would require a dose of 100-300 $\mu\text{g/kg}$, ten times higher than the intrathecal dose. Such intravenous analgesic doses are associated with undesirable sedative effects. On the other hand, the doses of 1-30 $\mu\text{g/kg}$, which produce analgesia at intrathecal doses, are known to have an anxiolytic, not sedative, effect when administered intravenously.

The spinal route of administration is especially useful because it avoids the side effects such as sedation and hypotension which are associated with the use of the above mentioned compounds systemically e.g. intravenously.

The present invention is illustrated by the following:

METHODS

Polyethylene catheters (8.5 cm) were inserted through the atlanto-occipital membrane with the tip at the L2 level to female Wistar rats (200-280 g) in intraperitoneal (i.p.) chloral hydrate anesthesia. After recovery for 3-5 days 400 μg of lidocaine was injected intrathecally (i.t.) and rats developing bilateral hind limb paralysis were accepted to the study. There were six rats in each group receiving either saline or dexmedetomidine in doses of 1.5, 3.0 or 6.0 μg in randomized order and double blind fashion in a volume of 10 μl . The tail-flick test (cut-off time 5 s) was performed before the i.t. injection and after 10, 20, 30, 45, 60, 90 min and 2, 3, 4, 5 and 6 h. In addition, in an open study, five rats were given atipamezole, a selective α_2 -adrenoceptor antagonist, 3 mg/kg i.p. before the injection of 6 μg of dexmedetomidine i.t.

RESULTS

To permit comparisons, the measured tail-flick latencies were converted to maximum percentage effect values (MPE) where $\text{MPE} = 100\% \times (\text{postinjection response latency} - \text{predrug response latency}) / (\text{cut-off time} - \text{predrug response latency})$. The mean maximum antinociceptive effect (91-99%) of dexmedetomidine in doses of 3 and 6 μg was reached within 10-20 min and the MPE's differed significantly from the control group for up to 5 h. The smallest dose of 1.5 μg caused a mean maximum effect of 48%, which lasted for 45 min. The MPE's stayed below 6% in the control group. Premedication with atipamezole abolished the antinociceptive effects of intrathecal dexmedetomidine.

The numerical data are summarized in Table 1. No signs of neurotoxicity into the spinal cord were detected.

It can be seen from this that compounds of the formula I, and in particular dexmedetomidine, causes pronounced dose dependent antinociceptive effect after doses of 1.5 to 6 $\mu\text{g/kg}$ per rat or 6 to 24 $\mu\text{g/kg}$. Thus the comparison of this dose to systemic effective doses in rats and humans suggests that the effective intrathecal doses in humans would be 0.05 to 0.5 $\mu\text{g/kg}$ compared to the effective systemic doses of 0.5-5 $\mu\text{g/kg}$.


These drugs could be administered to humans or other mammalian species as intrathecal or epidural injections or infusions to treat pain e.g. in surgical operations, cancer, spastic paraplegia or equivalent conditions. The injections or infusions may contain one or more diluents or carriers.

Table 1.

Mean \pm SEM values of the antinociceptive efficacy (%) at different time points after dexmedetomidine intrathecal administration.												
	Control (NaCl)			1.5 μ g DEX			3 μ g DEX			6 μ g DEX		
Time	n	mean	SEM	n	mean	SEM	n	mean	SEM	n	mean	SEM
10 min	6	-4.17	2.77	6	26.50	19.02	6	89.33	24.69	6	90.17	8.8
20 min	6	-6.67	2.69	6	35.00	18.92	6	98.67	1.33	6	91.00	9.00
30 min	6	-5.67	2.85	6	48.00	20.45	6	96.00	4.00	6	91.50	8.50
45 min	6	-5.17	1.69	6	45.00	17.98	6	95.33	4.67	6	91.00	9.00
60 min	6	-2.83	2.50	6	38.83	17.66	6	68.50	14.26	6	88.33	11.67
90 min	6	-2.83	2.06	6	31.83	17.02	6	62.50	18.37	6	86.00	9.66
2 h	6	-4.00	1.97	6	28.83	15.52	6	57.17	19.21	6	79.50	11.77
3 h	6	-2.33	1.82	6	12.33	12.07	6	53.33	19.01	6	76.67	13.72
4 h	6	-5.17	2.27	6	8.33	10.46	6	46.00	14.81	6	64.83	17.92
5 h	6	-3.17	1.49	6	8.50	7.82	6	38.33	13.03	6	66.50	13.97
6 h	6	-0.67	1.63	6	3.83	7.28	6	27.50	15.56	6	44.17	16.66
7 h	3	3.00	3.00	5	10.40	14.30	6	33.67	15.95	5	34.00	11.48
8 h	2	-3.00	3.00	2	8.50	8.50	4	39.00	20.52	2	51.50	32.50
24 h	6	-1.67	1.99	6	3.33	4.11	6	2.33	1.86	6	16.67	13.05

Claims

1. The use of a compound which is an imidazole of the formula (I):



where X is H or CH₃ and R₁ and R₂, which can be the same or different, are each H or CH₃, a stereoisomer thereof or a non-toxic, pharmaceutically acceptable salt thereof in the manufacture of a medicament for local epidural or intraspinal administration.

2. The use according to claim 1 in which the medicament is for administration in an amount corresponding to administration of 0.05 to 0.5 μ g/kg of imidazole of formula (I).

3. The use according to claim 1 or 2 in which the medicament is an analgesic.

4. The use according to any one of claims 1 to 3 where the compound is medetomidine or (\pm)-4-[1-(2,3-dimethylphenyl)ethyl]-1H-imidazole.

5. The use according to any one of claims 1 to 3 where the compound is dexmedetomidine or (+)-4-[1-(2,3-dimethylphenyl)ethyl]-1H-imidazole.

6. The use according to any one of claims 1 to 3 where the compound is detomidine or 4-(2,3-dimethylbenzyl)-1H-imidazole.



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 31 1245

DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
D,Y	EP-A-0 331 374 (FARMOS-YHTYMA OY) * Page 2, paragraphs 1,2; claims * -----	1-6	A 61 K 31/415
Y	ACTA ANASTESIOLOGICA BELGICA, vol. 40, no. 4, 1989, pages 147-261; T.F. MEERT et al.: "Effects of adrenaline, an alpha2-adrenoceptor agonist, the volume of injection, and the global pain state of the animal on the activity of epidural sufentanil" * Summary * -----	1-6	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			A 61 K
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		27 November 90	BERTE M.J.
CATEGORY OF CITED DOCUMENTS			
X: particularly relevant if taken alone		E: earlier patent document, but published on, or after the filing date	
Y: particularly relevant if combined with another document of the same category		D: document cited in the application	
A: technological background		L: document cited for other reasons	
O: non-written disclosure		-----	
P: intermediate document		&: member of the same patent family, corresponding document	
T: theory or principle underlying the invention			

